# Mycofix<sup>®</sup> Select 3.0

Naturally ahead in mycotoxin risk management!






With 4 active modules Mycofix® Select protects animals from mycotoxicoses



Synergistic Blend of Minerals<sup>1</sup>

<sup>1</sup>Authorized in EU, Regulation No 1060/2013



Biomin<sup>®</sup> BBSH<sup>®</sup> 797<sup>2</sup> <sup>2</sup>Authorized in EU, Regulation No 1016/2013 and 2017/930



**Phytogenic Substances** 

**Phycophytic Constituents** 

### **Recommended for:**

Rations, when trichothecenes are of main concern

- Pullets and laying hens
- Chicks in the starter phase
- Broiler, turkey and ducks
- Finisher pigs
- Beef cattle



- Integral solution to mycotoxin problems through the process of selective adsorption of mycotoxins with polar functional groups and microbial inactivation of trichothecenes.
- Reduced liver and intestinal tract membrane damages caused by the presence of mycotoxins in the feed.
- Reduces lesions and inflammations caused by trichothecenes.
- Stimulates the immune system to compensate the suppression caused by the presence of mycotoxins in the feed.





## Mycofix<sup>®</sup> Select 3.0

Naturally ahead in mycotoxin risk management!

#### Administration:

Mycofix® Select 3.0 is added directly into the feed during mixing.

#### Dosage:

Prevention:	0.5 – 1 kg/mt of finished feed
Pullets and laying hens:	1.5 – 2.5 kg/mt of finished feed*
Chicks in the starter phase:	1.5 – 2.5 kg/mt of finished feed*
Broiler, turkeys and ducks:	1.5 – 2.5 kg/mt of finished feed*
Finisher pigs	1.5 – 2.5 kg/mt of finished feed*
Beef cattle	10 – 50 g/cow/day*

\* depending on the contamination level and production phase

### **Packing:**

25 kg plastic bag in corrugated cartons

#### **Stability:**

18 months from production date

#### Storage:

Store in a dry place and avoid direct sunlight

resentation or on for business of information s. All trademarks of thirds, unless of the second states of the seco

Disclaimer: dsm-firmenich has used diligent care to ensure that the information provided herein is accurate and up-to-date, however, dsm-firmenich makes no representation or warranty, either expressly or implied, of the accuracy, reliability, or completeness thereof. The information provided herein contains scientific and product information for business to business use and does not constitute or provide scientific or medical advice, diagnosis, or recommendation for treatment. Country or region-specific information should be considered when labeling or advertising to the final consumer. In no event shall dsm-firmenich be liable for any damages arising from or reliance upon, or use of, any information provided herein. The content of this document is subject to change without further notice. Please contact your local dsm-firmenich representative for further details. All trademarks listed in this document are either (registered) trademarks of, or trademarks licensed by, the dsm-firmenich group of companies in the Netherlands and/or other countries, unless explicitly stated otherwise.